

The Longitudinal and Transverse responses in inclusive electron scattering within the RPA framework.

E. Bauer¹, A. Polls² and A. Ramos²

¹Departamento de Física, Facultad de Ciencias Exactas,
Universidad Nacional de La Plata,
La Plata, 1900, Argentina.

²Departament d'Estructura i Constituents de la Matèria,
Universitat de Barcelona,
Diagonal 647, E-08028, Spain.

February 8, 2008

Abstract

The longitudinal and transverse nuclear responses to inclusive electron scattering reactions are analyzed within the Random Phase Approximation (RPA) framework. Several residual interactions are considered and it is shown that the exchange terms in the RPA make very difficult to find an effective residual interaction capable of reproducing simultaneously the quasielastic peak of both the longitudinal and transverse responses. By means of a simplified model it is illustrated that the residual interaction used in a ring approximation must fulfill some restrictions in order to qualitatively reproduce the full RPA results.

PACS number: 21.65, 25.30.Fj, 21.60.Jz.

Keywords: Nuclear Electron Scattering.

In this work we present a study of the longitudinal and transverse nuclear response functions to inclusive electron scattering reactions within the Random Phase Approximation (RPA) framework in non-relativistic nuclear matter. It is only recently that two different procedures to evaluate the nuclear responses in the RPA scheme are available for a general finite range effective interaction [1, 2]. In the past, most of the calculations have been performed in the so-called Ring Approximation (RA). In fact, both the RA and RPA approximations account for the excitations of particle-hole type that can be induced by the electromagnetic probe but, formally, the RA is only the direct contribution of the RPA. It has been common to adjust the effective nucleon-nucleon interaction to reproduce the nuclear response in the quasielastic peak within the RA. We note that within the RA different pieces of the interaction act either in the longitudinal or in the transverse response, facilitating the extraction of the parameters of the interaction for a certain momentum transfer. Since the explicit evaluation of exchange terms is important, as we will see, one could aim to apply the same procedure in the framework of the RPA.

To find a force that gives a reasonable account of the basic features of both responses would be particularly interesting to define a proper framework beyond which other excitations, like ground state correlations, final state interactions and meson exchange currents, which have been shown to be important (see [3] and Refs. therein), are included. In the present brief report we have explored several representative interactions commonly used in the literature and, as we will show, no one is able to give a simultaneous satisfactory description of both longitudinal and transverse responses at various momentum transfer values. The reason lies in the fact that, as opposed to the RA, the full interaction is present in the RPA-exchange terms and in addition these terms are very important. Therefore the adjustment of one channel affects the other one and this means that reproducing both channels, longitudinal and transverse, is a far more complex task.

On the other hand, it has been common to use a modified residual interaction in the RA scheme trying to account for exchange terms. Another purpose of the present work is to show, making use of a simplified model, that while the RA is able to keep the qualitative trends of the complete RPA response, it is unable to provide a quantitative account of the exchange terms tied to the finite range part of the force.

As it will be shown below, the qualitative agreement can only be achieved if some simple constraints over the parameters of the interaction are observed.

The longitudinal and transverse pieces of the RPA response are calculated following the scheme of Ref. [1], but using a HF basis according to the prescriptions given in [4], where the effect of the HF self-energy is adjusted by means of a set of two effective masses (one for particles and the other one for holes) and an energy shift. Each set is chosen to reproduce the exact HF response and depends only on the momentum transferred by the electron. The RPA calculation is performed for three different interactions. The first one, labelled V^I , is described in [5] and contains the exchange of the mesons π , ρ and ω plus a g' term.

The structure of the second one has been widely used in the literature and consists of contact terms plus a $(\pi + \rho)$ -meson exchange interaction given by,

$$V^{II}(k) = \frac{f_\pi^2}{\mu_\pi^2} (f_0 + f'_0 \boldsymbol{\tau} \cdot \boldsymbol{\tau}' + g_0 \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + g'_0 \boldsymbol{\tau} \cdot \boldsymbol{\tau}' \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + V_\pi(k) + V_\rho(k)) \quad (1)$$

with

$$V_\pi(k) = -\Gamma_\pi^2(k) \frac{k^2}{k^2 + \mu_\pi^2} \boldsymbol{\sigma} \cdot \hat{\mathbf{k}} \boldsymbol{\sigma}' \cdot \hat{\mathbf{k}} \boldsymbol{\tau} \cdot \boldsymbol{\tau}', \quad (2)$$

$$V_\rho(k) = -C_\rho \Gamma_\rho^2(k) \frac{k^2}{k^2 + \mu_\rho^2} (\boldsymbol{\sigma} \times \hat{\mathbf{k}}) \cdot (\boldsymbol{\sigma}' \times \hat{\mathbf{k}}) \boldsymbol{\tau} \cdot \boldsymbol{\tau}', \quad (3)$$

where μ_π (μ_ρ) is the pion (rho) rest mass and $C_\rho = 2.3$. For the form factor of the πNN (ρNN) vertex we have taken

$$\Gamma_{\pi,\rho}(k) = \frac{\Lambda_{\pi,\rho}^2 - \mu_{\pi,\rho}^2}{\Lambda_{\pi,\rho}^2 + k^2}, \quad (4)$$

with $\Lambda_\pi = 1.3$ GeV and $\Lambda_\rho = 1.75$ GeV. The momentum transferred by the interaction is k and the static limit of the $(\pi + \rho)$ -meson exchange interaction has been taken. The parameters f_0, f'_0, g_0, g'_0 of this force have been left free so as to reproduce both the longitudinal and transverse responses. However, the presence of all terms of the interaction in the exchange contributions makes this purpose very difficult. This will be illustrated more clearly in the simplified model discussed below. An acceptable reproduction of both channels, especially the longitudinal one, for momentum transfer $q = 410$ MeV/c is found with the values $f_0 = -0.1$, $f'_0 = 0.2$, $g_0 = 0$ and $g'_0 = 0.5$.

Finally, we consider the parameterization of the Bonn potential of Ref. [4]. This interaction, which will be referred as V^{III} , is expressed as the exchange of π , ρ , σ and ω mesons.

For each interaction, the effective masses and energy shifts necessary to calculate the HF responses are shown in Table I.

In Figs. 1 and 2, we present the RPA results (solid lines) for a momentum transfer of $q = 300$ MeV/c and $q = 410$ MeV/c, respectively, and for the three interaction models. Our results are compared with the experimental data on ^{40}Ca [6]. Also we present the HF (long-dashed lines) and the RA (short-dashed lines) responses. The latter is obtained from the “contact” interaction model explained below. In all the calculations the Fermi momentum is taken to be $k_F = 235$ MeV/c, which is an appropriate value to simulate the results in finite nuclei [7]. The HF response is in fact qualitatively similar to the free Lindhard function but it is hardened by the presence of the single particle spectrum, which increases the energy of particle-hole excitations.

The behavior of the RPA responses in the longitudinal and transverse channels depends on the interaction considered. For the V^I and V^{II} interactions, when the position of the RPA peak for one channel is moved towards higher energies with respect to the peak in the HF response, the peak in the other channel moves in the opposite direction. This particular feature is a consequence of the exchange terms in the RPA. Only the V^{III} interaction gives a hardening of the response in both channels. We also observe that the interaction V^{II} , whose parameters have been adjusted to account for the response at $q = 410$ MeV/c, produces a response at $q = 300$ MeV/c in poor agreement with the experimental results. This means that the interaction should have a richer momentum dependence than just the one provided by π - and ρ -meson exchange.

In order to clarify this behavior, let us consider the simpler case of an interaction:

$$V_C = \frac{f_\pi^2}{\mu_\pi^2} (f + f' \boldsymbol{\tau} \cdot \boldsymbol{\tau}' + g \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + g' \boldsymbol{\tau} \cdot \boldsymbol{\tau}' \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + \\ + h \boldsymbol{\sigma} \cdot \hat{\mathbf{q}} \boldsymbol{\sigma}' \cdot \hat{\mathbf{q}} + h' \boldsymbol{\sigma} \cdot \hat{\mathbf{q}} \boldsymbol{\sigma}' \cdot \hat{\mathbf{q}} \boldsymbol{\tau} \cdot \boldsymbol{\tau}') \quad (5)$$

where f , f' , g , g' , h and h' are all constants. We have preferred to use the spin longitudinal terms h and h' instead of the tensor terms (proportional to S_{12}). This is because, at variance with S_{12} , the spin longitudinal terms do not interfere with the vector terms g and g' in the RA.

This interaction is now used to evaluate direct and exchange terms in the RPA. For such an interaction it is possible to account for exchange terms by a re-definition

of these constants as follows

$$\begin{aligned}
f_{ant} &= f - (f + 3f' + 3g + 9g' + h + 3h')/4 \\
f'_{ant} &= f' - (f - f' + 3g - 3g' + h - h')/4 \\
g_{ant} &= g - (f + 3f' - g - 3g' - h - 3h')/4 \\
g'_{ant} &= g' - (f - f' - g + g' - h + h')/4 \\
h_{ant} &= h - (h + 3h')/2 \\
h'_{ant} &= h' - (h - h')/2.
\end{aligned} \tag{6}$$

The first term in the *r.h.s.* of each equation is the direct contribution, while the terms between parenthesis come from the action of the exchange operator over the interaction. It is important to keep in mind that solving the RA with this new set of parameters is equivalent to solving the RPA with the original contact interaction of Eq. (5). At variance with the direct case, only three of the parameters f_{ant} - h'_{ant} , are independent. They are related through the following relations:

$$\begin{aligned}
g'_{ant} &= -(f_{ant} + h'_{ant})/3 \\
g_{ant} &= -(2f_{ant} - h'_{ant})/3 - f'_{ant} \\
h_{ant} &= -h'_{ant},
\end{aligned} \tag{7}$$

which are obtained by simply solving the system (6). Note that the spin longitudinal terms are usually assumed to be proportional to the momentum transfer and they cancel in the Landau limit. Formally, Eq. (5) with $h = h' = 0$, can be viewed as the zeroth order in the Legendre expansion of the Landau-Migdal interaction. In this sense, (7) is a particular case of the more general sum rules results of Ref. [8].

Within the RA, f_{ant} and f'_{ant} govern with the same weight the longitudinal response, while for the transverse one only g'_{ant} is relevant (g_{ant} gives also a contribution, but of the order $\mu_s^2/\mu_v^2 \approx 0.035$ with respect to that of g'_{ant}).

Now we turn back to the analysis of the RPA response with a general finite range interaction. For direct RPA terms, the external momentum fixes the momentum of the particle-hole interaction. That means that, for direct RPA terms, the interaction behaves as a contact one for each momentum transfer. In Table II we have extracted the “contact” terms of the interactions V^{I-III} for $q = 300$ MeV/c and $q = 410$

MeV/c. Working as if these parameters belonged to a contact interaction we have obtained, using Eqs. (6), the corresponding values $f_{ant}-h'_{ant}$, also shown in Table II. We have then performed a RA calculation with these values and the results are given by the short-dashed lines in Figs. 1-2. This simple model illustrates that if we represent a general interaction by a contact one by ignoring the momentum dependence of the finite range terms, the corresponding RPA (which is in fact a RA with the parameters redefined according to Eqs. (6)) gives a qualitative agreement with the result obtained using the complete interaction, which is displayed by the solid lines in Figs. 1-2. The position and shape of the quasielastic peak are qualitatively similar, in both longitudinal and transverse channels. Therefore, the momentum dependence of the finite range force in the exchange terms is not strong enough to modify the trends on the longitudinal and transverse responses induced by the simplified contact interaction. This, in turn, implies that the coefficients of the different terms of any effective interaction that might be used to account effectively for exchange terms in RA should obey to a high degree the restrictions imposed by the relations (7).

We can summarize our results by saying that the momentum dependence of the interaction in the exchange terms gives rise to quantitative differences between the RPA and the RA responses, the later one being calculated with the antisymmetrized contact version of the original force. This makes advisable to use the RPA with the complete interaction before attempting more complicated studies of other types of correlations. Correlations beyond RPA are certainly necessary since, to the best of our knowledge, no work is able to reproduce the quasi-elastic response at any momentum transfer for both longitudinal and transverse channels, using the same interaction for medium and heavy nuclei. We have unsuccessfully attempted to find one interaction that at least reproduces the position of the peaks for both channels within the RPA framework.

Another observation of the present work is that if one still wants to account for exchange terms within a simpler RA approach one should make sure that the effective interaction is such that the relations (7) are preserved to a high degree, since only in this case the RPA and RA are in qualitative agreement. The implementation of these relations is as simple as the RA itself. We consider significant this partial saving of the RA because of the extreme simplicity of the ring propagator in nuclear matter,

which contrasts with the difficult evaluation of the full RPA.

It is also important to note that, while it is possible to find an interaction that adjusts the longitudinal and transverse channels in a RA scheme, it will not obey the relations (7) and, therefore, the corresponding original direct interaction cannot be recovered. Such type of interactions cannot be blindly used in calculations of other types of many-body correlations that explicitly require the knowledge of the original interaction. In particular, one should avoid replacing the effective interaction by dressing it with the ring propagator if relations (7) are ignored.

References

- [1] E. Bauer, A. Ramos and A. Polls, Phys. Rev. **C54** (1996) 2959.
- [2] A. De Pace, Nucl. Phys. **A635** (1998) 163.
- [3] S. Boffi, C. Giusti, F. D. Pacati and M. Radici; Electromagnetic Response of Atomic Nuclei, Clarendon Press, Oxford 1996.
- [4] M. B. Barbaro, A. De Pace, T. W. Donnelly and A. Molinari, Nucl. Phys. **A596** (1996) 553.
- [5] R. Cenni and P. Saracco, Phys. Rev. **C50** (1994) 1851.
- [6] C. F. Williamson *et al.*, Phys. Rev. **C56** (1998) 3152.
- [7] J. E. Amaro, A. M. Lallena and G. C3, Int. J. Mod. Phys. **E3** (1994) 735.
- [8] B. L. Friman and A. K. Dhar, Phys. Lett. **85B** (1979) 1.
- [9] E. Bauer and A. Lallena, Phys. Rev. **C59** (1999) 2603.

Figure Captions

Figure 1: Response function for ^{40}Ca at momentum transfer $q = 300 \text{ MeV}/c$. The panels on the *l.h.s.* show the longitudinal response and those on the *r.h.s.* show the transverse one. Results are shown for the HF (long dashed line), the RA (short-dashed line) and the RPA responses (solid line). Results in rows (a), (b) and (c) are obtained with the V^I , V^{II} and V^{III} interactions, respectively. The experimental data was taken from [6].

Figure 2: The same as Fig. 1, but for a momentum transfer $q = 410 \text{ MeV}/c$.

Table I.

$q = 300 \text{ MeV/c}$			
int.	m^*/m		$\Delta\omega$ [MeV]
	particles	holes	
V^I	0.97	0.85	6.
V^{II}	0.83	0.67	7.
V^{III}	0.75	0.65	9.
$q = 410 \text{ MeV/c}$			
int.	m^*/m		$\Delta\omega$ [MeV]
	particles	holes	
V^I	1.0	0.85	7.
V^{II}	0.85	0.67	9.
V^{III}	0.78	0.65	11.

Table I: Effective masses and energy shifts for the different residual interactions described in the text and for two values of the momentum transfer. The energy shift is denoted by $\Delta\omega$.

Table II.

$q = 300 \text{ MeV/c}$						
int.	f	f'	g	g'	h	h'
V^I	0.000	0.000	-0.094	0.375	0.094	-0.465
V^{II}	-0.100	0.200	0.000	0.351	0.000	-0.578
V^{III}	-0.630	0.085	-0.085	-0.163	0.085	-0.570
	f_{ant}	f'_{ant}	g_{ant}	g'_{ant}	h_{ant}	h'_{ant}
V^I	-0.448	0.212	-0.162	0.397	0.744	-0.744
V^{II}	-0.581	0.394	-0.295	0.483	0.868	-0.868
V^{III}	0.300	0.042	-0.541	0.199	0.898	-0.898
$q = 410 \text{ MeV/c}$						
int.	f	f'	g	g'	h	h'
V^I	0.000	0.000	-0.148	0.282	0.148	-0.386
V^{II}	-0.100	0.200	0.000	0.267	0.000	-0.493
V^{III}	-0.442	0.041	-0.134	-0.251	0.134	-0.482
	f_{ant}	f'_{ant}	g_{ant}	g'_{ant}	h_{ant}	h'_{ant}
V^I	-0.270	0.189	-0.226	0.308	0.653	-0.653
V^{II}	-0.455	0.352	-0.295	0.398	0.739	-0.739
V^{III}	0.630	-0.080	-0.603	0.053	0.789	-0.789

Table II: For a fixed momentum the interaction for direct RPA terms is a constant. Columns f - h' are the values of these constants for the interactions V^{I-III} described in the text. Columns f_{ant} - h'_{ant} result from applying Eqs. (6) using the values of columns f - h' as direct contributions.




Fig. 1




Fig. 2